arXiv:1210.4687v3 [cond-mat.mes-hall] 27 Oct 2012

Prediction of Measurable Casimir-like Force and Free Energy Signatures of Exciton Condensates in DQWs

T. Hakioğlu$^{(\mathbf{1,2})}$, Ege Özgün$^{(\mathbf{1})}$, Mehmet Günay$^{(\mathbf{1})}$
(**1**) *Department of Physics, Bilkent University, 06800 Ankara, Turkey*
(**2**) *Institute of Theoretical and Applied Physics, 48740 Turunç, Muğla, Turkey*

We predict three measurable free energy related signatures of the condensed excitons in DQWs and propose experiments for their measurements. The first two are related to the static as well as dynamical measurements of the attractive critical Casimir force in the DQW geometry arising due to the presence of a sharp phase boundary defined by the critical coupling constant (i.e. the layer separation), the temperature, the exciton number and the electron-hole number imbalance. We also propose, in a somewhat similar geometry, a third experiment that distinguishes electrostatically the dependence of the free energy on the exciton concentration between a free exciton gas and an exciton condensate. The proposed experiments are regarded as highly conclusive on the presence of the exciton condensate, with a result irrespective from the relative strengths of the dark/bright components, a clear diffculty in understanding the nature of the condensed excitonic ground state.

PACS numbers: 71.35.-y,71.70.Ej,03.75.Hh,03.75.Mn

In the late 40's Casimir predicted the presence of an unusual attractive force between two uncharged metallic plates held in vacuum[1]. The Casimir force (CF) is a pure quantum mechanical effect manifested by the vacuum fluctuations of the electromagnetic field in the presence of metallic boundaries. The CF is attractive between the two infinite metallic plates and the force per unit area (the Casimir pressure) is given by $P_c = \mathcal{F}_c/A = -\pi^2\hbar c/(240 D^4)$ where A is the area of the plates and D is the separation between them. For typical values $A \simeq 1\mu m^2$ and $D \simeq 100\mathring{A}$, $\mathcal{F}_c \simeq -1.3\times 10^{-7}N$. The earliest successful measurements of the CF date back to ten years ago where a perfect agreement with the theory was achieved.[2].

One way to derive the CF is to calculate the difference between the vacuum free energies of a system with and without the boundaries. On the other hand, the idea of the vacuum fluctuations can be extended to a more general context including for instance, in confined geometries, the collective behaviour in the many particle macroscopically coherent ground states at low temperatures when the free energy becomes a function of the geometry. Here the macroscopically ordered ground state replaces the role of the vacuum. The role of the fluctuations around the vacuum are then played by the fluctuations around the critical point where a phase transition from the *normal* to macroscopically ordered ground state occurs. Therefore it is expected that some statistical mechanical systems at the boundary of a phase transition (critical region) also exhibit a similar effect, i.e. the critical Casimir force (CCF). Indeed, the CCF has been predicted[3] for binary liquid mixtures and measured[4]. The CCF, was also theoretically predicted for a BEC[5] but experimentally has not been yet observed. On the other hand, the Casimir-Polder force between a BEC and a semiconductor plane has been observed[6].

In the two dimensional coupled electron-hole systems confined to separate quantum wells with a layer separation on the order of an exciton Bohr radius a_B, excitons are expected to condense, below a T_c, into a phase coherent ground state. The free energy of the exciton condensate (EC) is a function of the pairing strength driven by the Coulomb interaction and, in contrast to the other statistical systems discussed above, the interaction strength is an exponential function of the layer separation D given by $V_{eh}(\mathbf{q}) = -(2\pi e^2/\varepsilon q)\, e^{-q\,D}$. Here $q = |\mathbf{q}|$ is the exchange momentum between the electrons and the holes and ϵ is the dielectric constant of the medium. In these systems, the condensate's order parameter, hence the free energy, is a sharp function of the layer separation in the vicinity of the phase boundary between the normal exciton gas and the EC.[7] The Casimir-like force that we theoretically predict in the excitonic DQWs is a new type of a Casimir effect which emerges at the phase boundary but is also strong deep inside the condensed phase.

In ECs one has a large number of controllable parameters like the total number of condensed excitons n_x, the electron-hole imbalance n_-, the layer separation D and the temperature T. The sharp phase boundary, separating the normal exciton gas from the EC, is defined by the critical parameters $n_x^{(c)}, n_-^{(c)}, D^{(c)}$ and T_c, which can be expressed by a critical manifold $T_c = g(D^{(c)}, n_x^{(c)}, n_-^{(c)})$. It was predicted that[7], in EC the phase boundary is present even at zero temperatures, i.e. $g(D^{(c)}, n_x^{(c)}, n_-^{(c)}) = 0$ where the free energy has a sharp second order phase transition across the critical manifold.

In our earlier numerical calculations based on the Hartree-Fock self consistent mean field approach[7] we observed this sharp phase boundary with a discontinuous derivative of the free energy with respect to D at the critical manifold. In this letter, we propose an analytical explanation for this effect. We also predict an order of magnitude of the CCF with its dependence on the critical parameters.

The phase boundary is determined by the condensate's free energy given by $\Delta\Omega = \Omega_{con} - \Omega_N \leq 0$, where Ω_{con} Ω_N are the total free energies in the condensed and the normal phase respectively. In this system we observe three types of dependence on D. The trivial dependence on D arises from the renormalized single particle energies that depend on the layer separation through multiple Coulomb processes between the electrons and the holes. The second is the critical fluctuations of the condensate around its mean value at T_c. The third dependence on D arises from the condensate's gap $\Delta = (\Delta_D^2 + \Delta_B^2)^{1/2}$ where Δ_D and Δ_B are the dark and the bright components of the order parameter[7, 8]. In this letter, we use the mean-field self consistent Hartree-Fock scheme at zero temperature which ignores the first two smooth dependences on D in comparison with the stronger electron-electron and hole-hole interactions. The total CCF than arises from the dependence of the condensation free energy $\Delta\Omega$ on Δ and is given by

$$\mathcal{F}_C = -\frac{\partial \Delta\Omega(T, V, \mu_x, \mu_-)}{\partial \Delta}\frac{\partial \Delta}{\partial D} \tag{1}$$

including all dependences on Δ. Here the free energy $\Omega = -k_B T \ln Tr\{exp(-\beta(\hat{H} - \mu\hat{N})\}$ is given up to a constant term by

$$\Omega_{con} = \sum_{\sigma,\mathbf{k}} \Big\{\sigma f_\sigma \frac{Tr\{\Delta\Delta^\dagger\}}{2E_\mathbf{k}} - \frac{2}{\beta}\ln\Big[1 + e^{-\beta(\epsilon_\mathbf{k}^{(-)} + \sigma E_\mathbf{k})}\Big]\Big\} \tag{2}$$

Here $\sigma = (+,-)$ are the spin states, $\Delta(\mathbf{k})$ is a 2×2 matrix order parameter describing the spin dependent condensate[8] including the dark and the bright components[9] with $Tr\{\Delta\Delta^\dagger\} = \sum_{\sigma,\sigma'} |\Delta_{\sigma\sigma'}|^2$, $f_\sigma = 1/[1 + e^{\beta(\epsilon_\mathbf{k}^{(-)} + \sigma E_\mathbf{k})}]$, $\epsilon_\mathbf{k}^{(-)} = (\xi_\mathbf{k}^{(e)} - \xi_\mathbf{k}^{(h)})/2$ and $\xi_\mathbf{k}^{(e)} = \hbar^2\mathbf{k}^2/(2m_e) - \mu_e + \Sigma_\mathbf{k}^{(e)}$, $\xi_\mathbf{k}^{(h)} = \hbar^2\mathbf{k}^2/(2m_h) - \mu_h + \Sigma_\mathbf{k}^{(h)}$ parameterized by the electron and the hole band masses m_e, m_h and the chemical potentials μ_e, μ_h, and $E_\mathbf{k} = [\epsilon_\mathbf{k}^2 + Tr\{\Delta\Delta^\dagger\}/2]^{1/2}$ where $\epsilon_\mathbf{k} = (\xi_\mathbf{k}^{(e)} + \xi_\mathbf{k}^{(h)})/2$. In what follows, we ignore the effect of the self energies $\Sigma_\mathbf{k}^{(e)}, \Sigma_\mathbf{k}^{(h)}$, assume that the electron and the hole masses are equal, whereas allow an imbalance between their concentrations[10]. We have then, $\epsilon_\mathbf{k}^{(-)} = -\mu_-$ where $\mu_- = (\mu_e - \mu_h)/2$.

At zero temperature, the order parameter is the solution of a self consistent equation given by

$$\Delta_\mathbf{k} = -\frac{\pi e^2}{\varepsilon}\int \frac{d\mathbf{q}}{(2\pi)^2}\frac{e^{-q\,D}}{q} G_{\mathbf{k}+\mathbf{q}} \tag{3}$$

where

$$G_\mathbf{k} = \frac{\Delta_\mathbf{k}\, F(\epsilon_\mathbf{k}^{(-)}, E_\mathbf{k})}{\sqrt{(\epsilon_\mathbf{k} - \mu_x)^2 + \Delta_\mathbf{k}^2}}\,. \tag{4}$$

Here $F(\epsilon_k^{(-)}, E_k) = \Theta(-\epsilon_k^{(-)} - E_k) - \Theta(-\epsilon_k^{(-)} + E_k)$. The self consistent scheme is completed with the constraints on the exciton number n_x and the electron-hole imbalance n_- yielding the exciton chemical potential $\mu_x = (\mu_e + \mu_h)/2$ and μ_-.

Analytic calculations here are more demanding than those in the constant interaction models in atomic condensates by the presence of the momentum dependent interaction $V_{eh}(\mathbf{q})$. One therefore resorts to numerical calculations requiring the self-consistent solution of the order parameter together with the constraints on the conserved number of particles. However, in certain regions in the momentum space one can understand the behaviour of the system analytically by using proper approximations which can restore the behaviour of the numerical solution around the phase boundary. In the large D limit, the strongest contribution in Eq. (3) comes in the region $\mathbf{q} \simeq 0$. Expanding $G_{\mathbf{k}+\mathbf{q}}$ upto second order in q, i.e. $G_{\mathbf{k}+\mathbf{q}} \simeq G_\mathbf{k} + \nabla_\mathbf{k} G_\mathbf{k}.\mathbf{q} + G''_\mathbf{k} q^2/2$ where $G''_\mathbf{k}$ is the second derivative with respect to k_x (or equivalently to k_y) at $\mathbf{k} = 0$ and examining Eq. (3) in the vicinity of $\mathbf{k} = 0$ by a similar parabolic approximation for $\Delta_\mathbf{k}$ where the first order terms in both expansions are not present due to the angular symmetry, we have a self consistency condition for Δ_0 and $\Delta''_0 = \Delta''_\mathbf{k}|_{\mathbf{k}=\mathbf{0}}$ given by

$$\Delta_0 = -E_0\{G_0 + \frac{2}{D^2}G''_0\}, \qquad E_0 = \frac{e^2}{2\epsilon D}$$
$$\Delta''_0 = -\frac{\Delta_0}{\mu_x}\frac{\hbar^2}{m} \tag{5}$$

We notice that Eq. (5) is easy to solve at the zero temperature since

$$\lim_{k\to 0} F(\epsilon_k^{(-)}, E_k)\Big|_{T=0} = \begin{cases} 1 & \text{for} \quad \Delta_0^2 + \mu_x^2 < \mu_-^2 \\ -1 & \text{for} \quad \Delta_0^2 + \mu_x^2 > \mu_-^2 \end{cases} \tag{6}$$

The first case in Eq. (6) is allowed when there is a high electron hole imbalance, indicated by a sufficiently large μ_-. For such large μ_- there is no non-zero solution for Δ_k in the limit $k \to 0$. If μ_- is zero or weak, a nonzero solution is allowed in the lower case which is independent of μ_-. Under these assumptions Δ_0 is found to be

$$\Delta_0 = \sqrt{E_0^2 - \mu_x^2} \tag{7}$$

with Δ''_0 as given by Eq. (5). As a result of this parabolic expansion, we have that

$$E_k = \frac{E_0}{\mu_x}|\epsilon_k - \mu_x| \tag{8}$$

We can also find approximate analytical relations for the exciton chemical potential μ_x and the relative chemical potential μ_- using the conservation of the electron n_e and the hole n_h charges with $n_x = (n_e + n_h)/2$ and $n_- = n_e - n_h$,

$$n_x = \frac{1}{2A}\sum_{\mathbf{k},\sigma}\Big\{(1 + \frac{\epsilon_\mathbf{k}^{(x)}}{E_\mathbf{k}})(2 + \sigma f_\sigma) + (1 - \frac{\epsilon_\mathbf{k}^{(x)}}{E_\mathbf{k}})(2 - \sigma f_\sigma)\Big\}$$
$$n_- = \frac{1}{A}\sum_{\mathbf{k},\sigma}(f_\sigma - \frac{1}{2})\,. \tag{9}$$

We find that, for $\mu_- \ll E_0$,

$$\mu_- = \frac{E_0}{2\Gamma\mu_x}\, n_- \tag{10}$$

$$\mu_x = -\Big(\frac{E_0}{2} - \frac{|n_-|}{8\Gamma}\Big) + \sqrt{\Big(\frac{E_0}{2} - \frac{|n_-|}{8\Gamma}\Big)^2 + \frac{E_0 n_x}{\Gamma}}$$

where $\Gamma = m_x/(2\pi\hbar^2)$ is the two-dimensional density of states with $m_x = (m_e^{-1} + m_h^{-1})^{-1}$ is the exciton reduced mass. Eq.'s (10) yield the chemical potentials $\mu_-(n_x, n_-)$ and $\mu_x(n_x, n_-)$, and inserting these into Eq.(7) we find $\Delta_0(E_0, n_x, n_-)$ where $E_0 \propto 1/D$. The result of this simple analytical approximation for Δ_0, μ_x as a function of n_-, n_x and D as well as the sharp phase boundary defined by the onset of $\Delta_0 = 0$ captures the main features of the numerical calculations in [7] for $|n_-| \ll n_x$.

This analytical approach also helps understanding the dependence of the critical Casimir force on E_0, n_x, n_- and D. We can use Eq.'s(7) and (10) to estimate the condensation energy $\Delta\Omega$ in Eq. (2) on the phase boundary. Since there is a weak dependence on n_-, we consider $n_- = 0$ and hence $\mu_- = 0$ in our estimate. Calculating Eq. (2) for this case, we find the simple result

$$\Delta\Omega = -\frac{\Gamma}{2\,E_0}[\mu_x(\mu_x^2 + 3\Delta_0^2) - \mu_0^3] \tag{11}$$

where $\mu_0 = n_x/2\Gamma$ is the chemical potential at the phase boundary given by $\Delta_0 = 0$. Using Eq.'s(7) and (10) in (11) and calculating Eq.(1) we find that at the phase boundary, the CCF is given by,

$$\frac{\mathcal{F}_C}{A} = -\frac{n_x^2}{2\Gamma D_c} \tag{12}$$

where A is the sample area and D_c is the critical layer separation at the phase boundary. The latter can be found from Eq. (7) for $\Delta_0 = 0$ as $D_c = e^2\Gamma/(\epsilon n_x)$. For a sample size of $A \simeq 10^3 \mu m^2$ and a typical concentration of $n_x \simeq 10^{11} cm^{-2}$, we find that $\mathcal{F}_C \simeq 10^{-9} N$. This analytical result is more reliable then our previous calculation [7] due to the fact that, the uncertainty in the numerical derivative can be very large at the phase boundary.

Due to the dielectric between the quantum wells, the direct measurement of $\mathcal{F}_C$ is much more challenging than measuring the electromagnetic CF between the two metallic plates separated by vacuum. To overcome this, we propose the experimental set-up in Fig.(1) based on detecting the CCF induced motion of a micro cantilever. By means of modulating n_x when the other parameters, i.e. n_-, D, are fixed, the $\mathcal{F}_c$ serves as a driving force for exciting static displacement as well as dynamical oscillations of the cantilever. Such mechanical oscillators can be made with high quality factors which have been successfully tested in MEMS recently.[12]

For the static experiment, the actuation of $\mathcal{F}_c$ is to deform tetragonally the lattice around the EC within the DQW. If the DQW is located asymmetrically in the beam with respect to the neutral plane, this will lead to bending of the beam in accordance with Fig.(1). A laser beam and an optical set-up to generate a specular reflection is placed on the bended structure in order to measure the deflection angle $\Delta\Theta$ as a function of n_x which can be controlled externally. Note that from Eq.(12) and the expression for D_c, $\Delta\Theta \propto n_x^3$ and any sharp dependence of the exciton concentration is attributed to the sharp change in the CCF. Using classical theory of beam bending under stress[13], we find that the angular deflection is given by $\Delta\Theta \simeq L_y^2\mathcal{F}_C/(12EI)$ where $L_y \simeq 100\mu m, L_x \simeq 10\mu m$ as indicated in Fig.(1), E is the Young's modulus and $I \simeq L_y W^3/3$ is the second area moment of the curved structure with $W = 0.3\mu m$ being the thickness of the cantilever. Using $E \simeq 80GPa$ in GaAs type materials, and for $F_c \simeq 10^{-9}N$, we find that $\Delta\theta \simeq 3\times10^{-4} rad$ which is possible to detect.[14] Alternatively, it is possible to detect the $\mathcal{F}_c$ driven mechanical oscillations of the cantilever. A typical cantilever oscillator[12] can be driven with a power consumption $P = m_{eff}\Omega_0^3\Delta z_{rms}^2/Q \simeq 2\times10^{-15}J/s$ where Δz_{rms} is the rms vibrational amplitude, $m_{eff} \simeq 10^{-10}kg$ is the effective mass of the cantilever, $\Omega_0/(2\pi) \simeq 20kHz$ is the resonance frequency, and $Q \simeq 2.5\times10^5$ is the quality factor of the resonator. In the Fig.(1) geometry at resonance, this leads to $\Delta z_{rms} \simeq 50nm$ which is detectable.



FIG. 1. (Color online) The proposed geometry for the static and dynamical measurements of the CCF when the DQW is grown (a) above, and (b) below the neutral plane of the cantilever.

We now propose a third method to directly probe the condensate's free energy by an electrostatic measurement as shown in Fig(2). In experiments, it is necessary to suppress the tunneling and the recombination of the electrons with the holes in the DQW structure. This is done by an E-field strength along the z-axis which is created by the potential difference V_{AB}. Experimentally, once the condensate is formed within the DQWs on the order of a few tens of ns, a potential difference V_{CD} is boosted up to generate the critical E-field strength E_c along the x-direction. The E_c is found by

$$\frac{1}{2}\epsilon E_c^2 = \frac{1}{A\,D}\,|\Delta\Omega| \tag{13}$$

which can be estimated using Eq.(11). We plot E_c in



FIG. 2. (Color online) The (AlGa)As DQW grown (thick red lines) with the undoped GaAs (as the dielectric tunneling barrier) is sandwiched between metallic-like plates formed by the semitransparent top gate A and the n^+GaAs bottom layer with Ohmic Au-terminal B. The E-field along the z-axis is provided by a bias potential V_{AB}. Along the lateral x direction the critical field E_c to break up the condensate is provided by the potential V_{CD} between the ohmic gold terminals C and D fabricated by lithographic recess process. The thickness t underneath the recess trench should be large enough to enable Ohmic contact formation without significant leakage between the lateral and vertical terminals.

Fig. (3) as a function of n_x. In the same figure, the second plot refers to the critical E-field strength for the excitonic gas, in which the incoherent pairs are formed with the free energy given by $\Delta\Omega_{inc} = -E_b n_x$ where $E_b \simeq 6meV$ is the binding energy of a single exciton. For $n_x \simeq 10^{10} cm^{-2}$ and $D \simeq a_B$ we find for the condensed case $E_c \simeq 40 - 50kV/cm$ compared to $4kV/cm$ that is required to break the incoherent pairs if they were not condensed. The current that is produced by breaking the EC or by breaking the incoherent excitons is read through the CD contacts. To expect a difference between the two cases, the calibration for the current is necessary which can be done by doing the experiment first above the critical temperature. In case of an EC the current produced should be a short and strong pulse, whereas in the case of an incoherent exciton gas, it is expected to be weaker, speading longer in time. The critical field strength E_c can be read at the onset of the current.



FIG. 3. (Color online) The E_c [scaled by $(2e_0/\epsilon a_B^3)^{1/2}$], with $e_0 = e^2/(2\epsilon a_B)$, versus n_x [scaled by $1/a_B^2$]. The plot (a) refers to the condensate and (b) is for the incoherent exciton gas. (The figure is generated for $D = a_B \simeq 100 \mathring{A}$ and $n_- = 0$.)

The experiments proposed here, due to their indiscrimination between the dark and the bright components, are complementary to the existing experiments in the literature. Due to the explicit dependence of the pairing interaction on the layer separation, the DQW EC is a natural candidate for the CCF experiments, which can be the first observation of the CCF in a low temperature condensate.

ACKNOWLEDGEMENTS

The authors are grateful to Klaus-Juergen Friedland (Paul-Drude Institute/Germany) for useful discussions concerning the experimental set-ups.

---

[1] H.B.G. Casimir, Proc. K. Ned. Akad. Wet. **51**, 793 (1948).
[2] G. Bressi, G. Carugno, R. Onofrio and G. Russo, Phys. Rev. Lett. **88**, 041804 (2002);
[3] M.E. Fisher and P.G. de Gennes, C.R. Acad. Sci. Paris, **B 287**, 207 (1978).
[4] C. Hertlein, L. Helden, A. Gambassi, S. Dietrich and C. Bechinger, Nature, **451**, 172 (2008); M. Fukuto, Y. F. Yano and P.S. Pershan, Phys. Rev. Lett. **94**, 135702 (2005); R. Garcia and M.H.W. Chan, Phys. Rev. Lett. **88**, 086101 (2002).
[5] Shyamal Biswas, J K Bhattacharjee, Dwipesh Majumder, Kush Saha and Nabajit Chakravarty, J. Phys. **43** (2010) 085305; P. A. Martin and V. A. Zagrebnov, Europhys. Lett. **73** 15 (2006).
[6] J. M. Obrecht, R. J. Wild, M. Antezza, L. P. Pitaevskii, S. Stingari and E. A. Cornell, Phys. Rev. Lett. **98**, 063201 (2007); D. M. Harber, J. M. Obrecht, J. M. McGuirk and E. A. Cornell, Phys. Rev. A **72** 033610 (2005).
[7] T. Hakioğlu and E. Özgün, Sol. Stat. Comm. **151**, 1045 (2011).
[8] M.A. Can and T. Hakioglu, Phys. Rev. Lett. **103** 086404 (2009); T. Hakioglu and M. Sahin, Phys. Rev. Lett. **98**, 166405 (2007).
[9] With s-wave like Coulomb interaction, assuming the presence of time reversal symmetry, which is assumed here, the dark components $\Delta_D = \Delta_{\uparrow\uparrow} = \Delta_{\downarrow\downarrow}$ and the bright component $\Delta_B = \Delta_{\uparrow\downarrow} = -\Delta_{\downarrow\uparrow}$.
[10] Including realistically the difference between the electron and the hole band masses is not crucial for the physics that follows.
[11] A. Siria, S. Huant, G. Auvert, F. Comin, and J. Chevrier, Nano. Res. Lett. **5** 1360 (2010).
[12] H. Yamaguchi, H. Okamoto, S. Ishihara, and Y. Hirayama, App. Phys. Lett. **100**, 012106 (2012).
[13] I.S. Sokolnikoff, Mathematical Theory of Elasticity, McGraw Hill, New York 1956.
[14] The resolution in $\Delta\Theta$ is expected to be similar to the radian accuracy of the conventional AFMs of $\simeq 10^{-5} rad$. [see Dawn A. Bonnell et al., Rev. Mod. Phys. **84**, 1343 (2012).]